

## Numerical modeling of air flows in an underground cavity connected to the surface by a shaft

Georges EDDE, Cerema - Grand Quevilly

Georges SADAKA, CNRS - Saint-Etienne du Rouvray

Raphaël ANTOINE, Cerema - Grand Quevilly

Ionut DANAILA, LMRS - Saint-Etienne du Rouvray

The Normandie region contains a lot of chalk cavities that were dug by farmers to fertilize fields. However, those cavities have a low dimension (2-3 meters height)/depth (20-40 meters depth) ratio which makes their detection very challenging with the classical geophysical methods.

Due to global warming, there will be more precipitations leading to the collapse of the cavities, causing the death of humans, thus their detection is a priority.

The Cerema center in France is testing an alternative method based on thermal infrared observations by drone<sup>1</sup>. So the aim of our model is to derive naturally thermal convection models in an underground cavity starting from available models at LMRS<sup>†</sup> in order to quantify the hydrothermal process at play in those cavities to compare it with the results of the drone. These numerical codes using finite elements make it possible to solve the Navier-Stokes equations and the heat equation in a fluid medium and a porous medium in 2D and 3D with a condition of solar radiation at the surface. Numerical simulations were carried out with the free software FreeFem++<sup>2</sup>.

The Figure 1 below shows the result of the simulation where we can see the triggering of natural air convection within and underground cavity opened to the atmosphere with an intensification of the airflow velocity with time and decreases with the downstream distance, respectively, and a number of cells of convection that change with time and the Rayleigh number. And so by comparing our result of the temperature to the one obtained with the drone, we can test if our model can be used to see the drone capability of detecting these underground cavities.




FIGURE 1 – Air circulation at  $t = 50$ s for natural convection simulated in the Barcq cavity (27170 - Eure - France) 3D with Rayleigh =  $1.e7$ .

Contact : [georges.edde@cerema.fr](mailto:georges.edde@cerema.fr)

1. C. Fauchard et al., *Guide technique- Laboratoire central des ponts et chaussées.*, (2004).
2. F. Hecht et al., *FreeFem++*, *Journal of Numerical Mathematics* 20 :251–266 (2012).