

Perfectly matched layers methods for mixed hyperbolic-dispersive equations

Maria Kazakova

Laboratoire de Mathématiques, Université Savoie Mont Blanc
collaboration with Christophe Besse, Sergey Gavrilyuk, Pascal Noble


31 Mai 2024

CANUM
Ile de Ré

Water waves models

$$\mu = H^2/L^2 \text{ (shallowness),}$$

$$\varepsilon = a/H \text{ (nonlinearity)}$$



$$\begin{cases} \frac{\partial h}{\partial t} + \nabla \cdot (h\mathbf{v}) = 0, & \text{(Mass Eq)} \\ \frac{\partial h\mathbf{v}}{\partial t} + \nabla \cdot \left(h\mathbf{v} \otimes \mathbf{v} + \frac{gh^2}{2}\mathcal{I} + p_{NH} \right) = 0, & \text{(Momentum Eq).} \end{cases}$$

model	NSWE $\mathcal{O}(\mu)$	$\mathcal{O}(\varepsilon\mu)$	SGN $\mathcal{O}(\mu^2)$
Pressure	$p_{NH} = 0$	Boussinesq	$p_{NH} = h^2\ddot{h}/3$
ε	no assump		no assump.
Type	hyperbolic		dispersive



Lannes, 2013

Water waves models

Hyperbolic vs Dispersif

Saint-Venant (**NSWE**)

Serre-Green-Naghdi (**SGN**)

Water waves models

Hyperbolic dispersive models

The most expensive step for non-hydrostatic models: elliptic problem

Recent advance on first-order hyperbolic equations with dispersive properties

📖 Favrie-Gavrilyuk, **2017** (SGN), Gavrilyuk et al. **2022** (BBM)

! Favrie-Gavrilyuk model is rigorously justified in 📖 Duchêne, **2019**


📖 Escalante et al. (artificial compressibility) **2019**

📖 Richard (compressible and quasi-incompressible) **2021**

Justificaton est en developpement (K. Msheik, V. Duchêne, A. Duran)

Boundary conditions


Problems are initially posed on infinite domain $x \in \mathbb{R}$



Boundary conditions

Problems are initially posed on infinite domain $x \in \mathbb{R} \rightarrow x \in \Omega$


Restriction of the observation area



Boundary conditions

Hyperbolic system - *Riemann-invariant form (if exist)*


Dispersive system - ?



Boundary conditions

Dispersive system - linear case: non-reflecting TBC, DTBC, PML

Nonlinear case (Coastal engineering, SGN) relaxation zones, sponge layers



Boundary condition

First strategy: Discrete Transparent BC for dispersive models


Dispersive systems, linear case:
Shrödinger (Ehrhardt, **2001**)
KdV, BBM (Besse et al., **2016**)
SGN (MK&Noble, **2020**)



MK, P.Noble (2020)

Boundary condition

Second strategy: Cartesian PML

Cartesian classical *Perfectly Matched Layers* (PML)  Bérenger (**1994**)

Boundary condition


Second strategy: Cartesian PML

Cartesian classical *Perfectly Matched Layers* (PML) 📖 Bérenger (1994)



Boundary condition


Second strategy: Cartesian PML

Cartesian classical *Perfectly Matched Layers* (PML)  Bérenger (1994)

Time domain \rightarrow frequency domain

PML change of variables


$$\mathbf{x} \in \mathbb{R}, \quad \tilde{\mathbf{x}} = \mathbf{x} \left(1 + \frac{\sigma(\mathbf{x})}{i\omega} \right)$$



in the layer $\sigma(x)$ linear functions, power functions or unbounded functions

Boundary condition


Second strategy: Cartesian PML

Cartesian classical *Perfectly Matched Layers* (PML)  Bérenger (1994)

Time domain \rightarrow frequency domain

PML change of variables


$$\mathbf{x} \in \mathbb{R}, \quad \tilde{\mathbf{x}} = \mathbf{x} \left(1 + \frac{\sigma(\mathbf{x})}{i\omega} \right)$$




in the layer $\sigma(x)$ linear functions, power functions or unbounded functions

$$\partial_{\tilde{x}} \rightarrow \left(1 + \frac{\sigma(x)}{i\omega} \right)^{-1} \partial_x$$

Well-posedness and stability


Following  Bécache *et al.*, **2003** ($\sigma(x) = \text{const}$) we can analyse stability

Well-posedness and stability

Following  Bécache *et al.*, **2003** ($\sigma(x) = \text{const}$) we can analyse stability
The initial equation admits *plane wave solution* of the form

$$u = Ue^{i(\mathbf{k}\cdot\mathbf{x}-\omega t)}, \quad \omega \in \mathbb{C}, \mathbf{k} \in \mathbb{R}$$

Well-posedness and stability


Following  Bécache *et al.*, **2003** ($\sigma(x) = \text{const}$) we can analyse stability
The initial equation admits *plane wave solution* of the form

$$u = U e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}, \quad \omega \in \mathbb{C}, \mathbf{k} \in \mathbb{R}$$

if and only if ω and \mathbf{k} are related via *dispersion relation*

$\mathfrak{F}(\omega, \mathbf{k}) = 0$, with solutions $\omega_j(\mathbf{k})$ which are called modes.

Well-posedness and stability

Following  Bécache *et al.*, **2003** ($\sigma(x) = \text{const}$) we can analyse stability
The initial equation admits *plane wave solution* of the form

$$u = U e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}, \quad \omega \in \mathbb{C}, \mathbf{k} \in \mathbb{R}$$

if and only if ω and \mathbf{k} are related via *dispersion relation*


$$\mathfrak{F}(\omega, \mathbf{k}) = 0, \text{ with solutions } \omega_j(\mathbf{k}) \text{ which are called modes.}$$

For the PML equation we define perturbed dispersion relation

$$\mathfrak{F}_{pml}(\omega, \mathbf{k}, \sigma) = 0, \text{ with modes } \tilde{\omega}_j(\mathbf{k}, \sigma)$$

$$\mathfrak{F}(\omega, \mathbf{k}) \rightarrow \mathfrak{F}_{pml}(\omega, \mathbf{k}, \sigma) \quad \text{with} \quad k \rightarrow k / \left(1 + \frac{i\sigma}{\omega}\right)$$

Well-posedness and stability

Following  Bécache *et al.*, **2003** ($\sigma(x) = \text{const}$) we can analyse stability
The initial equation admits *plane wave solution* of the form

$$u = U e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}, \quad \omega \in \mathbb{C}, \mathbf{k} \in \mathbb{R}$$

if and only if ω and \mathbf{k} are related via *dispersion relation*

$$\mathfrak{F}(\omega, \mathbf{k}) = 0, \text{ with solutions } \omega_j(\mathbf{k}) \text{ which are called modes.}$$

For the PML equation we define perturbed dispersion relation

$$\mathfrak{F}_{pml}(\omega, \mathbf{k}, \sigma) = 0, \text{ with modes } \tilde{\omega}_j(\mathbf{k}, \sigma)$$

$$\mathfrak{F}(\omega, \mathbf{k}) \rightarrow \mathfrak{F}_{pml}(\omega, \mathbf{k}, \sigma) \quad \text{with} \quad k \rightarrow k / \left(1 + \frac{i\sigma}{\omega}\right)$$

We search for solutions with an exponential behaviour and the PML equation is stable if and only if $\Im(\tilde{\omega}_j) \leq 0$ for all $\sigma \geq 0$.

Stability condition and inverse waves

We introduce notions of the *phase velocity* \mathbf{v}_p and the *group velocity* \mathbf{v}_g (general case $\mathbf{k} \in \mathbb{R}^3$):

Stability condition and inverse waves

We introduce notions of the *phase velocity* \mathbf{v}_p and the *group velocity* \mathbf{v}_g (general case $\mathbf{k} \in \mathbb{R}^3$):

$$\mathbf{v}_p(\mathbf{k}) = \frac{\omega_j(\mathbf{k})}{|\mathbf{k}|} \frac{\mathbf{k}}{|\mathbf{k}|}, \quad \mathbf{v}_g(\mathbf{k}) = \nabla_{\mathbf{k}} \omega_j(\mathbf{k}),$$
$$\forall \omega_j(\mathbf{k}) \text{ solution of } \mathfrak{F}(\omega, \mathbf{k}) = 0.$$

Stability condition and inverse waves

We introduce notions of the *phase velocity* \mathbf{v}_p and the *group velocity* \mathbf{v}_g (general case $\mathbf{k} \in \mathbb{R}^3$):

$$\mathbf{v}_p(\mathbf{k}) = \frac{\omega_j(\mathbf{k})}{|\mathbf{k}|} \frac{\mathbf{k}}{|\mathbf{k}|}, \quad \mathbf{v}_g(\mathbf{k}) = \nabla_{\mathbf{k}} \omega_j(\mathbf{k}),$$

$$\forall \omega_j(\mathbf{k}) \text{ solution of } \mathfrak{F}(\omega, \mathbf{k}) = 0.$$

Necessary stability conditions Bécache(2003)

If $\forall \mathbf{k} \in \mathbb{R}^3$, $(\mathbf{v}_p(\mathbf{k}) \cdot \mathbf{e}_j)(\mathbf{v}_g(\mathbf{k}) \cdot \mathbf{e}_j) \geq 0$,
the problem with classical Cartesian PML applied in \mathbf{e}_j direction is stable.

Stability condition and inverse waves


We introduce notions of the *phase velocity* \mathbf{v}_p and the *group velocity* \mathbf{v}_g (general case $\mathbf{k} \in \mathbb{R}^3$):

$$\mathbf{v}_p(\mathbf{k}) = \frac{\omega_j(\mathbf{k})}{|\mathbf{k}|} \frac{\mathbf{k}}{|\mathbf{k}|}, \quad \mathbf{v}_g(\mathbf{k}) = \nabla_{\mathbf{k}} \omega_j(\mathbf{k}),$$

$$\forall \omega_j(\mathbf{k}) \text{ solution of } \mathfrak{F}(\omega, \mathbf{k}) = 0.$$

Necessary stability conditions Bécache(2003)

If $\forall \mathbf{k} \in \mathbb{R}^3$, $(\mathbf{v}_p(\mathbf{k}) \cdot \mathbf{e}_j)(\mathbf{v}_g(\mathbf{k}) \cdot \mathbf{e}_j) \geq 0$,
the problem with classical Cartesian PML applied in \mathbf{e}_j direction is stable.



Stability condition and inverse waves

We introduce notions of the *phase velocity* \mathbf{v}_p and the *group velocity* \mathbf{v}_g (general case $\mathbf{k} \in \mathbb{R}^3$):

$$\mathbf{v}_p(\mathbf{k}) = \frac{\omega_j(\mathbf{k})}{|\mathbf{k}|} \frac{\mathbf{k}}{|\mathbf{k}|}, \quad \mathbf{v}_g(\mathbf{k}) = \nabla_{\mathbf{k}} \omega_j(\mathbf{k}),$$

$$\forall \omega_j(\mathbf{k}) \text{ solution of } \mathfrak{F}(\omega, \mathbf{k}) = 0.$$

Necessary stability conditions Bécache(2003)

If $\forall \mathbf{k} \in \mathbb{R}^3$, $(\mathbf{v}_p(\mathbf{k}) \cdot \mathbf{e}_j)(\mathbf{v}_g(\mathbf{k}) \cdot \mathbf{e}_j) \geq 0$,
the problem with classical Cartesian PML applied in \mathbf{e}_j direction is stable.

If there are backward propagating waves in the PML direction
the PML system is unstable.

Stability condition and inverse waves

We introduce notions of the *phase velocity* \mathbf{v}_p and the *group velocity* \mathbf{v}_g (general case $\mathbf{k} \in \mathbb{R}^3$):

$$\mathbf{v}_p(\mathbf{k}) = \frac{\omega_j(\mathbf{k})}{|\mathbf{k}|} \frac{\mathbf{k}}{|\mathbf{k}|}, \quad \mathbf{v}_g(\mathbf{k}) = \nabla_{\mathbf{k}} \omega_j(\mathbf{k}),$$

$$\forall \omega_j(\mathbf{k}) \text{ solution of } \mathfrak{F}(\omega, \mathbf{k}) = 0.$$

Necessary stability conditions Bécache(2003)

If $\forall \mathbf{k} \in \mathbb{R}^3$, $(\mathbf{v}_p(\mathbf{k}) \cdot \mathbf{e}_j)(\mathbf{v}_g(\mathbf{k}) \cdot \mathbf{e}_j) \geq 0$,
the problem with classical Cartesian PML applied in \mathbf{e}_j direction is stable.

Necessary stability conditions in the 1D case $v_g(k)v_p(k) \geq 0$

Cartesian PML: Typical exemple KdV

KdV equation

$$u_t + u u_x + \varepsilon u_{xxx} = 0, \quad \forall x \in \mathbb{R}, \quad \forall t > 0.$$

Cartesian PML: Typical exemple KdV

linear KdV equation

$$u_t + U u_x + \varepsilon u_{xxx} = 0 \quad \forall x \in \mathbb{R}, \quad \forall t > 0 \quad (TD)$$

Cartesian PML: Typical exemple KdV

In Frequency domain (after Fourier transform)

$$-i\omega\hat{u} + U\hat{u}_x + \varepsilon\hat{u}_{xxx} = 0 \quad \forall x \in \mathbb{R}$$

Cartesian PML: Typical exemple KdV

Artificial truncation by PML: $x \in \Omega, \quad \forall t > 0$

$$-i\omega\left(1 + \frac{i\sigma}{\omega}\right)u + U\partial_x u + \varepsilon\partial_x \left(\left(1 + \frac{i\sigma}{\omega}\right)^{-1}\partial_x \left(\left(1 + \frac{i\sigma}{\omega}\right)^{-1}\partial_x u \right) \right) = 0$$

Cartesian PML: Typical exemple KdV

$$-i\omega\left(1 + \frac{i\sigma}{\omega}\right)u + U\partial_x u + \varepsilon\partial_x \left(\left(1 + \frac{i\sigma}{\omega}\right)^{-1}\partial_x \left(\left(1 + \frac{i\sigma}{\omega}\right)^{-1}\partial_x u \right) \right) = 0$$

+ auxiliary variables u_1 and u_2 :

$$\partial_x u = \left(1 + \frac{i\sigma}{\omega}\right)u_1, \quad \partial_x u_1 = \left(1 + \frac{i\sigma}{\omega}\right)u_2,$$

Back to time domain

$$\partial_t u + \sigma u + U\partial_x u + \varepsilon\partial_x u_2 = 0,$$

$$\partial_t (u_1 - \partial_x u) + \sigma u_1 = 0, \quad \partial_t (u_2 - \partial_x u_1) + \sigma u_2 = 0.$$

$(TD)_{PML}$

By applying the initial value theorem, one finds

$$u_1|_{t=0} = \partial_x u|_{t=0}, \quad u_2|_{t=0} = \partial_{xx} u|_{t=0}.$$

The linear KdV equation

$$u_t + U u_x + \varepsilon u_{xxx} = 0, \quad \forall x \in \mathbb{R}, \quad \forall t > 0.$$

Proposition

- If $U = 0$, equations $(TD)_{PML}$ are always unstable.
- If $\varepsilon U < 0$, equations $(TD)_{PML}$ are stable if and only if $k^2 \geq 16 \frac{|U|}{|\varepsilon|}$.
- If $\varepsilon U > 0$, equations $(TD)_{PML}$ are stable if and only if $k^2 \leq \frac{U}{3\varepsilon}$.

Proof. The dispersion relation of $(TD)_{PML}$: Following  Bécache2003

$$\begin{aligned} &\text{dispersion relation for KdV with } k \rightarrow k/(1 + \frac{i\sigma}{\omega}) \\ &(\omega + i\sigma)^3 = kU(\omega + i\sigma)^2 - \varepsilon k^3 \omega^2. \end{aligned}$$

If $k = 0$, $\omega = -i\sigma$ and the condition $\Im(\omega) \leq 0$ is satisfied.

$$\text{If } k \neq 0 \quad \omega^2(\omega - \omega_0(k)) = 0, \quad \omega_0(k) = kU - \varepsilon k^3.$$

Two roots are bifurcating from 0 and one root bifurcates from $\omega = \omega_0(k)$.

The linear KdV equation

From straightforward computations, a necessary condition is

$$(U - \varepsilon k^2)(U - 3\varepsilon k^2) > 0$$

Here $v_g(k) = U - 3\varepsilon k^2$ and $v_p(k) = U - \varepsilon k^2$.

$$v_g(k)v_p(k) \geq 0.$$

So we recover the classical condition in the PML framework.

We have proved that $\Im(\omega) \leq 0$ for $\sigma > 0$ small enough, under conditions on k claimed in the proposition.

We show then that for any $\sigma > 0$, there are no real solutions, which means that $\Im(\omega) \neq 0$.

We conclude that these conditions are sufficient to guarantee stability, using continuity of the roots of a complex polynomial with respect to its coefficients. This ends the proof.

The linear KdV equation

Discretization

We consider a centered space FD with a Crank Nicolson in time scheme:

$$x_j = j\delta x, j \in \mathbb{Z}, t_n = n\delta t, n \in \mathbb{N}$$

$$2\frac{v_j^n - u_j^n}{\delta t} + \sigma v_j^n + U\frac{v_{j+1}^n - v_{j-1}^n}{2\delta x} + \varepsilon\frac{v_{2,j+1}^n - v_{2,j-1}^n}{2\delta x} = 0,$$

$$\frac{2}{\delta t} \left(\left(v_{1,j}^n - \frac{v_{j+1}^n - v_{j-1}^n}{2\delta x} \right) - \left(u_{1,j}^n - \frac{u_{j+1}^n - u_{j-1}^n}{2\delta x} \right) \right) + \sigma v_{1,j}^n = 0,$$

$$\frac{2}{\delta t} \left(\left(v_{2,j}^n - \frac{v_{1,j+1}^n - v_{1,j-1}^n}{2\delta x} \right) - \left(u_{2,j}^n - \frac{u_{1,j+1}^n - u_{1,j-1}^n}{2\delta x} \right) \right) + \sigma v_{2,j}^n = 0,$$

with $v_{k,j}^n = \frac{u_{k,j}^{n+1} + u_{k,j}^n}{2}$ for $k = 0, 1, 2$ and $u_{0,j}^n = u_j^n$.

The linear KdV equation

Numerical simulation

Case: $\varepsilon U > 0$





Initial condition $u_0(x) = \exp(-40(x+3)^2)$, $u_1 = u_0'$ and $u_2 = u_0''$.

The domain is $[-8, 8] \times [0, 200]$, $\delta x = 0.05$, $\delta t = \delta x$.

$$\sigma(x) = 2 \left(\max\left(0, \frac{x-5}{3}\right)^4 + \max\left(\frac{-x-5}{3}, 0\right)^4 \right)$$

$\varepsilon = U\delta x^2/4$ (stable case)

$\varepsilon = U\delta x^2/2$ (unstable case)



Representation of the function $v(t, x) = \log(1 + 1000|u(t, x)|)$.



The linear KdV equation

Case: $\varepsilon U < 0$

Initial condition $u_0(x) = \exp(-(x - 3)^2) \sin(2x)$.

$$\varepsilon = 16|U|\delta x^2$$

$$\varepsilon = 32|U|\delta x^2.$$



Representation of the function $v(t, x) = \log(1 + 1000|u(t, x)|)$



The linear KdV equation

Case: $\varepsilon U < 0$

Initial condition $u_0(x) = \exp(-(x - 3)^2) \sin(2x)$.

$$\varepsilon = 16|U|\delta x^2$$

$$\varepsilon = 32|U|\delta x^2.$$



Message I


We recover in this analysis the classical stability condition
 Since the phase and group velocities do not always have the same sign
 the PML for KdV is not always stable.

A hyperbolic KdV system

We now consider a relaxation of the original Korteweg-de Vries equation.

$$u_t + u u_x + \varepsilon \psi_x = 0, \quad p_t - \frac{p_x - \psi}{\tau} = 0, \quad \psi_t + \frac{u_x - p}{\tau} = 0,$$

ε – the dispersion parameter, $\tau > 0$ – the relaxation parameter.

Also: Euler-Lagrange equations for a given Lagrangian  [<hal>](#)

Formally, $\tau \rightarrow 0$, the function u turns out to be an approximate solution of the KdV equation. Indeed, p, ψ expand as

$$p = u_x + \tau u_{txx} + O(\tau^2), \quad \psi = u_{xx} + \tau (u_{txxx} - u_{tx}) + O(\tau^2).$$

By inserting this expansion we have

$$(u - \tau u_{xx} + \tau u_{xxxx})_t + u u_x + \varepsilon u_{xxx} = O(\tau^2).$$


which is the Benjamin-Bona-Mahoney (BBM) regularization of the KdV.




A hyperbolic KdV system

We now consider a relaxation of the original Korteweg-de Vries equation.

$$u_t + u u_x + \varepsilon \psi_x = 0, \quad p_t - \frac{p_x - \psi}{\tau} = 0, \quad \psi_t + \frac{u_x - p}{\tau} = 0,$$


ε – the dispersion parameter, $\tau > 0$ – the relaxation parameter.

Also: Euler-Lagrange equations for a given Lagrangian  [<hal>](#)



A hyperbolic KdV system

Initial wave: $u_0(x) = \exp(-40(x+2)^2)$



$v(t, x) = \log(1 + 1000|u(t, x)|)$ in the (x, t)
 in the case $\varepsilon U > 0$, $U = 1$, $\varepsilon = 5\delta x^2$
 (unstable for the original KdV)

Message II

PML are not always stable for hyperbolic approximation.


We can construct a stable version, but it will not be “exactly” PML

Application to abcd-model

We consider the hyperbolic-dispersive systems which models water wave propagation BBM-Boussinesq type model (also known as abcd-model):

abcd

$$\begin{aligned} (1 - b\partial_x^2)\partial_t\eta + \partial_x u + a\partial_x^3 u &= 0, \\ (1 - d\partial_x^2)\partial_t u + \partial_x \eta + c\partial_x^3 \eta &= 0, \end{aligned} \quad \forall (t, x) \in [0, T] \times [x_\ell, x_r].$$

 Bona, Chen and Saut (2002)


By-product: KdV dynamic is included in this model (properly chosed initial data creates approximate one-way propagating waves)

Application to abcd-model

We consider the hyperbolic-dispersive systems which models water wave propagation BBM-Boussinesq type model (also known as abcd-model):

abcd

$$\begin{aligned} (1 - b\partial_x^2)\partial_t\eta + \partial_x u + a\partial_x^3 u &= 0, \\ (1 - d\partial_x^2)\partial_t u + \partial_x \eta + c\partial_x^3 \eta &= 0, \end{aligned} \quad \forall (t, x) \in [0, T] \times [x_\ell, x_r].$$

 Bona, Chen and Saut (2002)

$$\begin{aligned} \partial_t(\eta - b\eta_2) + \sigma(\eta - b\eta_2) + \partial_x(u + au_2) &= 0, \\ \partial_t(u - du_2) + \sigma(u - du_2) + \partial_x(\eta + c\eta_2) &= 0, \\ \partial_t(\eta_1 - \partial_x \eta) + \sigma\eta_1 &= 0, \quad \partial_t(\eta_2 - \partial_x \eta_1) + \sigma\eta_2 = 0, \\ \partial_t(u_1 - \partial_x u) + \sigma u_1 &= 0, \quad \partial_t(u_2 - \partial_x u_1) + \sigma u_2 = 0. \end{aligned}$$

The initial conditions are given by

$$\eta_i|_{t=0} = \partial_x \eta_{i-1}|_{t=0}, \quad u_i|_{t=0} = \partial_x u_{i-1}|_{t=0}, \quad i = 1, 2.$$

Application to abcd-model

Necessary condition

Denote v_g and v_p respectively the group velocity and phase velocity. A necessary condition of stability is written again $v_g(k)v_p(k) \geq 0$ for all $k \in \mathbb{R}$.

Proposition

The PML equations associated to the classical Boussinesq equation ($a = b = c = 0, d > 0$) and the shallow water equations with surface tension ($a = b = d = 0, c < 0$) are stable.

Proposition

The PML system is stable under the assumption $a = d = 0$ and $b > 0, c < 0$. The PML system is also stable in the case $b = c = 0$ and $d > 0, a < 0$.

Application to abcd-model



Bidirectionnel wave propagation

$$\eta(t = 0, x) = \exp(-x^2), \quad u(t = 0, x) = 0.$$

In order to chose a right propagating wave we need to set:

$$u(t = 0, x) = (1 - d\partial_x^2)^{-1/2}\eta(t = 0, x).$$

The FFT and inverse FFT allow to calculate the fractional derivative.



Application to abcd-model

Bidirectionnel wave propagation

$$\eta(t = 0, x) = \exp(-x^2), \quad u(t = 0, x) = 0.$$

In order to chose a right propagating wave we need to set:

$$u(t = 0, x) = (1 - d\partial_x^2)^{-1/2}\eta(t = 0, x).$$

The FFT and inverse FFT allow to calculate the fractional derivative.

Message III

The PML is always stable when dispersive properties of the model are better suited for this technique, i.e. the condition $v_g(k)v_p(k) \geq 0$ is always satisfied.

Conclusions

Results on PML stability for linearised water wave problem:

- PML is not suitable for KdV, partially for the hyperbolic version: hyperbolization does not help.
- PML works for large class of BBM-Boussinesq equations
- DTBC are better when $v_g(k)v_p(k) < 0$ (which is a common situation in dispersive problems).

I. Dispersive properties of the model are important for stability of PML

II. If the dispersive properties of the model do not fits to the necessary stability condition

Chose another model

Construct a non-classical PML